

A PHASE 1 STUDY OF DAILY ORAL TRC102 (METHOXYAMINE) TO ENHANCE THE THERAPEUTIC EFFECTS OF PEMETREXED IN PATIENTS WITH ADVANCED REFRACTORY CANCER

SP Anthony, MS Gordon, LS Rosen, GJ Weiss, L Liu, Y Xu, BJ Adams, D Alvarez, SL Gerson, CP Theuer, BR Leigh. Evergreen Hematology & Oncology, Spokane WA; Premiere Oncology, Scottsdale AZ & Santa Monica CA; Scottsdale Clinical Research Institute/TGen, Scottsdale AZ; University Hospitals Ireland Cancer Center and Case Comprehensive Cancer Center, CWRU, Cleveland OH; TRACON Pharmaceuticals Inc, San Diego CA

Presentation Overview

- TRC102 Background
- Study Objectives
- Study Design
- Study Results
- Summary & Conclusions

TRC102 Background

- TRC102 (methoxyamine) is a small molecule inhibitor of base-excision repair (BER) that enhances the cytotoxicity of alkylator and antimetabolite chemotherapy and reverses chemotherapy resistance (Liu et al, 2002; Liu et al, 1999; Bulgar et al, 2006)
- TRC102 inhibits BER by binding apurinic (AP) sites produced by glycosylase removal of damaged DNA bases, thereby targeting them for cleavage by topoisomerase II and triggering apoptosis (Yan et al, 2007)

Study Objectives

- Evaluate the safety and tolerability of escalating doses of TRC102 in combination with pemetrexed in patients with advanced or metastatic solid cancers
- Evaluate pharmacokinetics, tumor response, and pharmacodynamics by AP site assay

Study Design

- Phase 1, first in human, open-label, dose escalation study conducted at 3 institutions in the United States
- Oral TRC102 was escalated in cohorts of 3-6 patients in combination with a standard dose of i.v. pemetrexed

Dosing Schedule

- All patients received TRC102 alone, dosed daily on Days 1-4 of an initial 2 week cycle, followed by the combination of pemetrexed on Day 1 and TRC102 on Days 1-4 every 3 weeks thereafter
- In Cycle 3, the Day 1 TRC102 dose was held in order to obtain the AP site assay sample for pemetrexed alone

	Cycle 1 (2 Weeks)	Cycle 2 (3 Weeks)	Cycle 3 (3 Weeks)	Cycle 4+ (3 Weeks)
Oral TRC102 Dosing	Days 1-4	Days 1-4	Days 2-4	Days 1-4
Pemetrexed Dosing	None	Day 1	Day 1	Day 1


Eligibility Criteria

- Key inclusion criteria:
 - Adults (age \geq 18 years) with advanced or metastatic solid cancer for whom curative therapy is unavailable
 - ECOG performance status of 0 or 1
 - Adequate organ function
- Key exclusion criteria:
 - Receipt of cancer treatment within 4 weeks of study start
 - History of primary or secondary brain tumors
 - Significant pericardial, pleural or peritoneal effusions

Patient Demographics

Characteristic	Number of Patients (n=12)
Median Age	69
Gender	Female: 8 Male: 4
Screening ECOG Performance Status	ECOG 0: 4 ECOG 1: 8
Number of Prior Regimens	Median: 2 Range: 1 to 6
Race	Caucasian: 10 Black/African American: 1 Hispanic/Latino: 1

Treatment Status



TRC102 Pharmacokinetics


Dose (mg/m ²)	Cycle	Day	N	C-max* (ng/mL)	Half-life* (hr)	AUC* (hr-ng/mL)
15	1	1	4	20.5 (12.7 - 57.4)	35.7 (24.8 - 41.2)	291 (219 - 1132)
15	1	4	3	68.1 (27.0 - 129)	41.5 (36.0 - 52.2)	1136 (188 - 2580)
15	2	1	3	21.4 (12.7 - 56.0)	35.8 (5.6 - 55.4)	258 (126 - 1186)
30	1	1	6	43.9 (16.9 - 205.0)	21.8 (15.9 - 43.6)	716 (263 - 3462)
30	1	4	5	115.0 (29.2 - 282.0)	30.9 (26.5 - 59.4)	1829 (585 - 5230)
30	2	1	6	56.2 (14.2 - 137.0)	22.3 (12.6 - 52.5)	894 (222 - 2071)

- PK Conclusions

- TRC102 plasma concentrations required for in vivo activity are achievable at 15 and 30 mg/m² with daily oral administration
- TRC102 exposure increased proportionally between 15 and 30 mg/m²
- TRC102 accumulated with daily dosing (Days 1-4), but did not accumulate between cycles

TRC102 Pharmacodynamics

- TRC102 inhibits BER by covalently binding to AP sites produced in response to pemetrexed therapy. Pemetrexed was expected to expose AP sites when dosed alone on Cycle 3 Day 1. During Cycle 2 Day 1, TRC102 was expected to bind AP sites, thereby preventing them from being detected in the AP site assay.
 - AP sites were formed in the absence of TRC102 therapy on Cycle 3 Day 1
 - AP sites were reduced when TRC102 was administered in combination with pemetrexed



TRC102 Safety

- No adverse events were attributed to TRC102 when dosed alone in Cycle 1
- One patient who received pemetrexed + 30 mg/m² TRC102 experienced Grade 3 anemia during Cycle 2 that was considered a dose-limiting toxicity, recovered after 2 units PRBCs and a 50% TRC102 dose-reduction, and remained on study through Cycle 5

Related Adverse Events

TRC102 Possibly Related Events			
Preferred Term	N=12 Treated Patients		
	Grade 1	Grade 2	Grade 3
Anemia		2	2 (1=DLT)
Leukopenia			1
Neutropenia			3
Abdominal pain lower	1		
Diarrhea	1		
Nausea	1		
Retching	1		
Asthenia	1		
Fatigue	1	1	
Edema peripheral	1		
Pain		1	
Pyrexia	1		
Cellulitis		1	
Anorexia	2		
Dehydration		1	
Hypomagnesemia	1		
Insomnia	1		
Petechiae	1		
Pruritus	1		
Purpura	1		
Rash	1		

Hematologic Adverse Events

All Hematologic Events				
Preferred Term	N=12 Treated Patients			
	Grade 1	Grade 2	Grade 3	Grade 4
Anemia	1	3	3 (1=DLT)	
Leukopenia			1	
Neutropenia			4	1
Thrombocytopenia			1	



TRC102 + Pemetrexed Efficacy

- Efficacy information is available on the first 9 patients including the following:
 - Ongoing partial response after 2 cycles of pemetrexed + 30 mg/m² TRC102 in a female patient with squamous cell cancer of the tonsil metastatic to the right lung and right hilum (prior therapies included radiation therapy, paclitaxel/cetuximab, and carboplatin/paclitaxel)
 - Mixed response after 2 cycles of pemetrexed + 30 mg/m² TRC102 in a female patient with epithelial clear cell cancer of the ovary metastatic to the liver, off study in Cycle 6 for rising CA-125 (prior therapies included carboplatin/paclitaxel, cisplatin, and liposomal doxorubicin)
 - Stable disease for 9 cycles of pemetrexed + 30 mg/m² TRC102 as 3rd line therapy for a male patient with metastatic squamous cell lung cancer

Partial Response – Hilar Mass

After 2 cycles of
pemetrexed plus
TRC102 at 30 mg/m²

Baseline



Summary & Conclusions

- Daily oral TRC102 was well-tolerated at doses that achieved plasma levels associated with in vivo activity
- Pemetrexed and TRC102 co-administration did not alter the PK of either compound
- TRC102 accumulated with daily dosing in a manner consistent with its half-life >24 hours, but did not accumulate between cycles
- Pharmacodynamic data are consistent with the TRC102 mechanism of action
- There is early evidence of activity of the combination of pemetrexed + TRC102, including one partial response in a patient with refractory metastatic head and neck cancer
- Phase 2 studies are planned in multiple indications